



40th ANNUAL

International Conference of the
IEEE Engineering in Medicine
and Biology Society

Honolulu, Hawaii

July 17 to 21, 2018

 **MEDLINE**
U.S. National Library of Medicine

 **PubMed**




HAWAII
CONVENTION CENTER
Where Business and Aloha Meet

PROGRAM



40th Annual International Conference of the IEEE Engineering in Medicine and Biology Society



July 17-21 2018

Hawaii Convention Center
Honolulu, Hawaii

Conference Chairs

James Weiland
Michelle Khine

Program Chairs

Greg Suaning
Olaf Doessel



Indexed in PubMed® and MEDLINE®,
Products of the United States National
Library of Medicine



- 17:15-19:00 WePoS-01.3
Tracking the Time Varying Neural Tuning via Adam on Point Process Observations
 Zitong, Zhang (*Zhejiang Univ.*); Chen, Shuhang* (*Hong Kong Univ. of Science and Technology*); Yang, Zaiyue (*Southern Univ. of Science and Technology*); Wang, Yiwen (*Hong Kong Univ. of Science and Technology*)
- 17:15-19:00 WePoS-01.4
Pipeline for Forward Modeling and Source Imaging of Magneto-cardiographic Recordings via Spatiotemporal Kalman Filtering
 Habboush, Nawar* (*University of Kiel*); Hamid, Laith (*University of Kiel*); Siniatchkin, Michael (*University of Kiel*); Stephani, Ulrich (*Christian-Albrechts-University of Kiel*); Galka, Andreas (*Christian-Albrechts-University of Kiel*)
-
- WePoS-02: 17:15-19:00 Exhibit Hall 2
Biosignal Processing for Motor Imagery Studies – Poster Session (Theme 1) (Poster Session)
- 17:15-19:00 WePoS-02.1
Comparison of Different EEG Signal Analysis Techniques for an Offline Lower Limb Motor Imagery Brain-Computer Interface
 Ortiz, Mario* (*Universidad Miguel Hernández*); Rodriguez-Ugarte, Marisol (*Miguel Hernández University of Elche*); Ianez, Eduardo (*Universidad Miguel Hernandez de Elche*); Azorin, Jose M. (*Universidad Miguel Hernandez de Elche*)
- 17:15-19:00 WePoS-02.2
Sparse Kernel Machines for Motor Imagery EEG Classification
 Oikonomou, Vangelis (*Centre for Research and Technology Hellas*); Nikolopoulos, Spiros (*Information Technologies Institute, Centre for Research and Technology Hellas*); Petrantonakis, Panagiotis* (*Information Technologies Institute, Centre for Research and Technology-Hellas (CERTH)*); Kompatsiaris, Ioannis (Yannis) (*Information Technologies Institute, CERTH*)
- 17:15-19:00 WePoS-02.3
Analysis and Classification for EEG Patterns of Force Motor Imagery using Movement Related Cortical Potentials
 Wang, Kun* (*Tianjin University*); Xu, Minpeng (*Tianjin University*); Zhang, Shanshan (*Tianjin University*); Ke, Yufeng (*Tianjin University*); Ming, Dong (*Tianjin University*)
- 17:15-19:00 WePoS-02.4
FBCSP-Based Multi-Class Motor Imagery Classification using BP and TDP Features
 Abbas, Waseem* (*Lahore University of Management Sciences*); Khan, Nadeem Ahmad (*Signal Image and Video Processing Lab, Electrical Engineering, Syed Babar Ali School of Science and Engineering, Lahore University*)
- 17:15-19:00 WePoS-02.5
DeepMI: Deep Learning for Multiclass Motor Imagery Classification
 Abbas, Waseem* (*Lahore University of Management Sciences*); Khan, Nadeem Ahmad (*Signal Image and Video Processing Lab, Electrical Engineering, Syed Babar Ali School of Science and Engineering, Lahore University*)
- 17:15-19:00 WePoS-02.6
Emotion Recognition for Brain Machine Interface: Non-linear Spectral Analysis of EEG Signals using Empirical Mode Decomposition
 Esmailbeigi, Hananeh (*Univ. of Illinois at Chicago (UIC)*); Carella, Tommaso (*Univ. of Illinois at Chicago*); De Silvestri, Matteo (*Univ. of Illinois at Chicago*); Finedore, Mary* (*Univ. of Illinois at Chicago*); Haniff, Isaac (*Univ. of Illinois at Chicago*)
- 17:15-19:00 WePoS-02.7
Unsupervised Phase Learning and Extraction from Repetitive Movements
 Jatesiklat, Prayook (*NTU*); Anopas, Dollaporn* (*Nanyang Technological University*); Ang, Wei Tech (*Nanyang Technological University*)
- 17:15-19:00 WePoS-02.8
EEG Processing to Discriminate Transitive-Intransitive Motor Imagery Tasks: Preliminary Evidences using Support Vector Machines
 Catrambone, Vincenzo* (*Univ. di Pisa*); Greco, Alberto (*Univ. of Pisa*); Averta, Giuseppe (*Univ. of Pisa*); Bianchi, Matteo (*Univ. of Pisa*); Vanello, Nicola (*Univ. of Pisa*); Bicchi, A. (*Univ. of Pisa*); Valenza, G. (*Univ. of Pisa*); Scilingo, E.P. (*Univ. of Pisa*)
- 17:15-19:00 WePoS-02.9
Evaluation of Different Signal Processing Methods in Time and Frequency Domain for Brain-Computer Interface Applications
 Arnin, Jetsada* (*University of Strathclyde*); Kahani, Danial (*University of Strathclyde*); Lakany, Heba (*University of Strathclyde*); Conway, Bernard A. (*University of Strathclyde*)
- 17:15-19:00 WePoS-02.10
Real-Time Human Physical Activity Recognition with Low Latency Prediction Feedback using Raw IMU Data
 Mascaret, Quentin* (*Laval University*); Biemann, Mathieu (*Laval University*); Cheikh Latyr, Fall (*Université Laval*); Bouyer, Laurent (*University of Laval*); Gosselin, Benoit (*Laval University*)
-
- WePoS-03: 17:15-19:00 Exhibit Hall 2
Connectivity, Causality and Phase Locking in Biomedical Signals – Poster Session (Theme 1) (Poster Session)
- 17:15-19:00 WePoS-03.1
EEG based Network Connectivity Classification in 7 and 9 Years-Old Children
 Almabruk, Tahani A. A.* (*Omar Al-Mukhtar University*); Tan, Tele (*Curtin University*); Khan, Masood Mehmood (*Curtin University*)
- 17:15-19:00 WePoS-03.2
Low Gamma Band Cortico-Muscular Coherence Inter-Hemisphere Difference following Chronic Stroke
 Bao, Shi-chun* (*Chinese Univ. of Hong Kong*); Wong, Wan-wa (*Chinese Univ. of Hong Kong*); Leung, Wai Hong (*Chinese Univ. of Hong Kong*); Tong, Kai Yu, Raymond (*Chinese Univ. of Hong Kong*)
- 17:15-19:00 WePoS-03.3
Preliminary Evaluation of Fetal Congenital Heart Defects Changes on Fetal-Maternal Heart Rate Coupling Strength
 Alangari, Haitham M.* (*Khalifa University*); Kimura, Yoshitaka (*Tohoku University*); Khandoker, Ahsan H (*Khalifa University of Science, Technology and Research*)
- 17:15-19:00 WePoS-03.4
Complex Modulation Method for Measuring Cross-Frequency Coupling of Neural Oscillations
 Malinowska, Urszula (*Johns Hopkins University School of Medicine*); Zieleniewska, Magdalena (*University of Warsaw*); Boatman-Reich, Dana (*Johns Hopkins School of Medicine*); Franaszczuk, Piotr* (*U.S. Army Research Lab*)
- 17:15-19:00 WePoS-03.5
Modulation of Low-Frequency Pulsed Magnetic Field on Hippocampal Neural Oscillation in Depression Rats
 Wang, Ling (*Tianjin University*); Yang, Jiajia* (*Tianjin University*); Wang, Faqi (*Tianjin University*); Zhou, Peng (*Tianjin University*); Wang, Kun (*Tianjin University*); Ming, Dong (*Tianjin University*)
- 17:15-19:00 WePoS-03.6
Assessment of EEG Connectivity Patterns in Mild Cognitive Impairment using Phase Slope Index
 Gomez, Carlos* (*University of Valladolid*); Ruiz, Saúl J. (*BioMedical Engineering Group, University of Valladolid*); Poza, Jesus (*University of Valladolid*); Maturana-Candelas, Aarón (*University of Valladolid*); Núñez, Pablo (*University of Valladolid*); Pinto, Nádia (*Institute of Molecular Pathology and Immunology of the University of Porto (IPATIMUP)*); Tola-Arribas, Miguel A. (*Dept. of Neurology, Hospital Universitario Río Hortega*); Cano, Mónica (*Dept. of Clinical Neurophysiology, Hospital Universitario Río Hortega*); Hornero, Roberto (*University of Valladolid*)

Assessment of EEG Connectivity Patterns in Mild Cognitive Impairment Using Phase Slope Index

Carlos Gómez, *Senior Member, IEEE*, Saúl J. Ruiz-Gómez, Jesús Poza, *Senior Member, IEEE*, Aarón Maturana-Candelas, Pablo Núñez, Nádía Pinto, Miguel A. Tola-Arribas, Mónica Cano, and Roberto Hornero, *Senior Member, IEEE*

Abstract— Mild cognitive impairment (MCI) is a pathology characterized by an abnormal cognitive state. MCI patients are considered to be at high risk for developing dementia. The aim of this study is to characterize the changes that MCI causes in the patterns of brain information flow. For this purpose, spontaneous EEG activity from 41 MCI patients and 37 healthy controls was analyzed by means of an effective connectivity measure: the phase slope index (*PSI*). Our results showed statistically significant decreases in *PSI* values mainly at delta and alpha frequency bands for MCI patients, compared to the control group. These abnormal patterns may be due to the structural changes in the brain suffered by patients: decreased hippocampal volume, atrophy of the medial temporal lobe, or loss of gray matter volume. This study suggests the usefulness of *PSI* to provide further insights into the underlying brain dynamics associated with MCI.

I. INTRODUCTION

Mild cognitive impairment (MCI) is a pathology characterized by a cognitive deficit higher than expected for a given age and educational level, but insufficient to meet the dementia criteria [1]. Although the cognitive activity of MCI patients is impaired, they maintain their independence in functional and social skills. Recent studies have estimated that the MCI prevalence is between 12 and 18% for subjects over 60 years [2]. Some of them remain stable over time, but more than 50% progress to different types of dementia such as dementia due to Alzheimer’s disease (AD) [3], vascular dementia [4], Parkinson disease [5], or dementia with Lewy bodies [6]. In particular, the rate of conversion to dementia

This research was supported by ‘Ministerio de Economía y Competitividad’ and ‘European Regional Development Fund’ under project TEC2014-53196-R, by ‘European Commission’ and ‘European Regional Development Fund’ under project ‘Análisis y correlación entre el genoma completo y la actividad cerebral para la ayuda en el diagnóstico de la enfermedad de Alzheimer’ (‘Cooperation Programme Interreg V-A Spain-Portugal POCTEP 2014-2020’), and by ‘Consejería de Educación de la Junta de Castilla y León’ under project VA037U16. P. Núñez and S. J. Ruiz are in receipt of predoctoral grants co-financed by the ‘Junta de Castilla y León’ and ESF. N. Pinto’s work is partially financed through the FCT postdoctoral grant SFRH/BPD/97414/2013 and projects POCI-01-0145-FEDER-007274 and UID/MAT/00144/2013.

C. Gómez, Saúl J. Ruiz-Gómez, J. Poza, A. Maturana-Candelas, P. Núñez, and R. Hornero are with the Biomedical Engineering Group, E.T.S. Ingenieros de Telecomunicación, University of Valladolid, Paseo Belén 15, 47011 Valladolid, Spain (e-mail: carlos.gomez@tel.uva.es).

N. Pinto is with the Institute of Molecular Pathology and Immunology of the University of Porto (IPATIMUP), the Institute for Research and Innovation in Health Sciences, and the Center of Mathematics of University of Porto, Portugal.

M. A. Tola is with the Department of Neurology, Hospital Universitario Río Hortega, Valladolid, Spain.

M. Cano is with the Department of Clinical Neurophysiology, Hospital Universitario Río Hortega, Valladolid, Spain.

due to AD is approximately 15% per year [3]. Additionally, the medication used to delay symptoms is more effective in the early stages of dementia. For both reasons, an AD early diagnosis is mandatory.

Electroencephalography (EEG) is a non-invasive technique that allows recording the electric fields generated by the brain. EEG is widely used due to its portability, availability and low cost. In addition, EEG has already shown its usefulness to characterize neuronal dynamics in MCI and AD [7,8].

Brain activity from MCI and AD patients has been widely analyzed with spectral techniques and nonlinear methods. Spectral techniques revealed a slowing of the EEG activity [9], whereas nonlinear methods showed a complexity decrease and a regularity increase of the spontaneous brain activity [10]. However, all these methods measure local activation at the sensor level without taking into account the possible interaction between the different EEG channels. Stam and van Straaten [11] suggested that simple activation studies are not enough to provide a complete characterization of brain dynamics. For this reason, connectivity analysis has focused a great attention in recent years. Previous studies showed lower connectivity values between EEG channels in patients with MCI using Pearson Correlation Coefficient, Magnitude Squared Coherence (*MSCOH*), and Phase Lag Index (*PLI*) [12,13]. All these measures estimate functional connectivity (i.e. the dependence between remote neurophysiological events). However, it is essential to know the directionality of the relationships to characterize completely the dynamics of the information flow in the brain network. To overcome these limitations, measures of effective or causal connectivity are needed (i.e. the influence that one neuronal system exerts on another), such as Granger Causality (*GC*), Transfer Entropy (*TE*), or Phase Slope Index (*PSI*). The advantage of *PSI* over other measures of effective connectivity, such as *GC* and *TE*, lies in its insensitive to false interactions due to volume conduction [14]. The aim of this work is to analyze the EEG connectivity patterns in MCI patients and in healthy controls (HC) by means of an effective connectivity measure (*PSI*).

II. MATERIALS AND METHODS

A. Subjects

Spontaneous EEG activity was analyzed in 41 MCI patients (17 men and 24 women, age = 76.1 ± 6.8 years, mean \pm standard deviation) and 37 HC subjects (12 men and 25 women, age = 76.3 ± 3.8 years). No significant differences between groups were found in age (p -value > 0.05 , Kruskal-Wallis test) nor in gender (p -value > 0.05 , Chi-square test).

TABLE I. SOCIO-DEMOGRAPHIC AND CLINICAL DATA FOR EACH GROUP

	HC group	MCI group
Number of subjects	37	41
Age (years) (Mean \pm SD)	76.3 \pm 3.8	76.1 \pm 6.8
Gender (Male:Female)	12:25	17:24
Education level (A:B) ^a	11:26	29:12
MMSE ^b (Mean \pm SD)	28.8 \pm 1.1	27.2 \pm 1.7
B-ADL ^c (Mean \pm SD)	1.2 \pm 0.2	3.0 \pm 0.9

^a A: primary education or below; B: secondary education or above

^b MMSE: Mini-Mental State Examination

^c B-ADL: Bayer-Activities of Daily Living

Mini-Mental State Examination test (MMSE) was used to evaluate the cognitive deficit of the subjects, while the functional status was assessed by means of the Bayer-Activities of Daily Living scale (B-ADL). None of the participants in the study was taking any medication that could influence on the EEG data. Table 1 shows the sociodemographic and clinical data for each group.

All participants and patients' caregivers were informed about the research background and the study protocol. Moreover, all of them gave their written informed consent to be included in the study. The Ethics Committee at the "Hospital Universitario Río Hortega" (Valladolid, Spain) endorsed the study protocol, according to The Code of Ethics of the World Medical Association (Declaration of Helsinki).

B. EEG recording

EEG activity was recorded with a 19-channel equipment (XLTEK[®], Natus Medical, Pleasanton, CA). The subjects were relaxed, awake and with their eyes closed during the EEG acquisition. EEG signals were acquired at a sampling frequency of 200 Hz for the following 19 electrodes placed following the International System 10-20: Fp1, Fp2, Fz, F3, F4, F7, F8, Cz, C3, C4, T3, T4, T5, T6, Pz, P3, P4, O1 and O2. Independent Component Analysis (ICA) was then applied to minimize the presence of oculographic, cardiographic and myographic artifacts. Afterwards, signals were digitally processed with a FIR filter with a Hamming window between 1 and 70 Hz and with a notch filter at 50 Hz. Finally, artifact-free epochs of 5-s length (1000 samples) were selected by visual inspection.

C. Phase Slope Index (PLI)

PSI is method proposed by Nolte *et al.* [14] that allows estimating the direction of the information flux. This measure is based on the slope of the phase of cross-spectra between two time series. The cross-spectra for two EEG channels $x_i(t)$ and $x_j(t)$ is defined as:

$$S_{ij}(j) = \langle \hat{x}_i(f) \hat{x}_j^*(f) \rangle, \quad (1)$$

where $\langle \cdot \rangle$ denotes expectation value and $\hat{x}_i(f)$ denotes the Fourier transform for the time series $x_i(t)$.

PSI is based on the delay associated to the cause-effect relationship between the times series corresponding to two EEG channels. Assuming that the interaction between these

two signals is merely a delay by time, $x_j(t)$ can be expressed as:

$$x_j(t) = cx_i(t - \tau), \quad (2)$$

where c is a constant. For the cross-spectra between the two EEG channels:

$$S_{ij}(f) = \langle \hat{x}_i(f) \hat{x}_j^*(f) \rangle \propto e^{i2\pi f\tau} \equiv e^{i\Phi(f)}. \quad (3)$$

Equation (3) shows that the phase-spectrum $\Phi(f) = 2\pi f\tau$ is linear and proportional to the time delay τ . Therefore, if the slope of $\Phi(f)$ is positive, the causal direction is estimated to go from $x_i(t)$ to $x_j(t)$. On the other hand, a negative slope of $\Phi(f)$ indicates that the information flow goes from $x_j(t)$ to $x_i(t)$.

Taking into account additional requirements, as statistical robustness and insensitivity to signals that do not interact, the *PSI* between two time series $x_i(t)$ and $x_j(t)$ can be defined as [14]:

$$\tilde{\Psi}_{ij} = \Im(\sum_{f \in F} Coh_{ij}^*(f) Coh(f + \delta f)), \quad (4)$$

where $Coh_{ij}(f) = S_{ij}(f) / \sqrt{S_{ii}(f) S_{jj}(f)}$ is the complex coherency, $S_{ij}(f)$ is the cross-spectral matrix, δf is the frequency resolution at a given frequency band F , and $\Im(\cdot)$ denotes the imaginary part.

Finally, it is convenient to normalize $\tilde{\Psi}$ by an estimate of its standard deviation:

$$\Psi = \tilde{\Psi} / \text{std}(\tilde{\Psi}), \quad (5)$$

with $\text{std}(\tilde{\Psi})$ being estimated using jackknife method [14]. *PSI* values around zero must be ignored in order to improve the robustness and reliability of the method [15].

III. RESULTS

PSI results were calculated at the EEG typical frequency bands: δ (1-4 Hz), θ (4-8 Hz), α (8-13 Hz), β_1 (13-19 Hz), β_2 (19-30 Hz), and γ (30-70 Hz). Figure 1 shows the mean *PSI* values for MCI and HC groups at the connections that showed statistically significant differences between both groups. The third column of Figure 1 illustrates the corresponding *p*-values (Mann-Whitney *U* test): red color tones indicate significant increases in connectivity in MCI patients compared to HC subjects, while blue color tones indicate significant decreases in MCI group in comparison with HC one. Our results revealed that EEG activity in MCI patients is characterized by a decrease in effective connectivity, mainly at δ and α bands. Specifically, *PSI* values show a significant decrease of the information flow from the lateral brain regions to the frontal area in δ and γ bands, from the frontal area to the posterior one in the θ band, and from central and posterior regions to the left lateral area in δ and α frequency bands.

Receiver operating characteristic (ROC) curves with a leave-one-out cross-validation procedure were used to assess the ability of *PSI* to discriminate MCI patients from HC subjects. The highest accuracy value was obtained for the connection T6 to Fp2 at δ band: 71.79% (sensitivity of 78.38%, specificity of 65.85%, and area under the ROC curve of 0.71).




Figure 1. Connectivity patterns obtained with *PSI* for HC (left column) and MCI (central column) groups, where the arrows indicate the direction of the information flow. Right column shows the statistically significant differences between groups: red color tones indicate significant increases in connectivity in MCI patients compared to HC subjects, while blue color tones indicate decreases. Connections were only displayed when statistically significant differences between groups were found (p -values < 0.05 , Mann-Whitney U test).

IV. DISCUSSION AND CONCLUSIONS

In this study, spontaneous EEG activity was analyzed from 41 MCI patients and 37 HC subjects using *PSI*, a measure of effective connectivity. Our results show that *PSI* values are significantly lower for patients with MCI at the low frequency bands. Causal measures have been previously applied to EEG and functional magnetic resonance imaging (fMRI) data, revealing diminished connectivity between different brain areas [16,17]. Additionally, our *PSI* results showed an alteration of the information flow patterns in all the frequency bands, being more remarkable for long distance connections and for low frequency bands. These results agree with previous studies that reported a connectivity decrease between the frontal region and the lateral and posterior areas in the δ and θ bands using functional measures, such as *PLI* [12]. Causal measures, as *GC* and Directed Transfer Function (*DTF*), also revealed a connectivity decrease between the frontal and posterior areas, although the affected frequency bands differ [13,18]. This inconsistency on the results may be due to the heterogeneity of MCI, to the different neuroimaging techniques, to the different connectivity coupling measures, or to a combination of these factors. Clinically, the observed abnormal patterns may be due to the structural brain changes suffered by patients with MCI: decreased hippocampal volume, atrophy of the medial temporal lobe, or loss of gray matter volume [19].

Finally, an accuracy of 71.79% was achieved using a ROC curve with a leave-one-out cross-validation procedure. Previous studies reported accuracies between 69.8% and 83% for HC vs. MCI classification using different coupling measures [13,16,20]. For instance, accuracy values of 69.8% were reached when *MSCOH* and synchronization likelihood were applied to magnetoencephalographic (MEG) recordings [20]. Dauwels *et al.* [16] yielded classification rates of 83% using *DTF*. Finally, *GC* method correctly classified 79.5% of the MCI and HC subjects who participated in a recent study.

It is important to mention some limitations of this research work. First, it would be convenient to increase the sample size to augment the statistical power of the results. Second, we have analyzed all the possible combinations between the different EEG electrodes in the current study; it would be interesting to divide the brain into regions to try to facilitate the interpretation of the results in spite of losing spatial resolution. Finally, it would be interesting to use other measures of effective connectivity, such as *GC*, *DTF*, or *TE*, to characterize brain information flow patterns from different perspectives.

We can conclude that the analysis of EEG activity using *PSI* method revealed that MCI causes significant changes in the brain activity. This study constitutes an initial step to demonstrate the possible application of novel measures of effective connectivity, such as *PSI*, for the characterization of AD in early stages.

REFERENCES

- [1] B. Winblad, K. Palmer, M. Kivipelto, V. Jelic, L. Fratiglioni, L.O. Wahlund, *et al.*, "Mild cognitive impairment - Beyond controversies, towards a consensus: Report of the International Working Group on Mild Cognitive Impairment," *J. Intern. Med.*, vol. 256, pp. 240–246, 2004.
- [2] RC. Petersen, "Mild cognitive impairment," *Contin. (Minneapolis)*, vol. 22, pp. 404–418, 2016.
- [3] C. Davatzikos, P. Bhatt, L.M. Shaw, K.N. Batmanghelich, and J.Q. Trojanowski, "Prediction of MCI to AD conversion, via MRI, CSF biomarkers, and pattern classification," *Neurobiol. Aging*, vol. 32, pp. 1–17, 2011.
- [4] J.S. Meyer, G. Xu, J. Thornby, M.H. Chowdhury, and M. Quach, "Is mild cognitive impairment prodromal for vascular dementia like Alzheimer's disease?," *Stroke*, vol. 33, pp. 1981–1985, 2002.
- [5] C.C. Janvin, J.P. Larsen, D. Aarsland, and K. Hugdahl, "Subtypes of mild cognitive impairment in Parkinson's disease: Progression to dementia," *Mov. Disord.*, vol. 21, pp. 1343–1349, 2006.
- [6] J. Molano, B. Boeve, T. Ferman, G. Smith, J. Parisi, D. Dickson, *et al.*, "Mild cognitive impairment associated with limbic and neocortical Lewy body disease: A clinicopathological study," *Brain*, vol. 133, pp. 540–556, 2010.
- [7] J. Dauwels, F. Vialatte, and A. Cichocki, "Diagnosis of Alzheimer's disease from EEG signals: where are we standing?," *Curr. Alzheimer Res.*, vol. 7, pp. 487–505, 2010.
- [8] J. Jeong, "EEG dynamics in patients with Alzheimer's disease," *Clin. Neurophysiol.*, vol. 115, pp. 1490–1505, 2004.
- [9] M. Baker, K. Akrofi, R. Schiffer, and M.W. O'Boyle, "EEG patterns in mild cognitive impairment patients," *Open Neuroimag. J.*, vol. 2, pp. 52–55, 2008.
- [10] C.J. Stam, "Nonlinear dynamical analysis of EEG and MEG: Review of an emerging field," *Clin. Neurophysiol.*, vol. 116, pp. 2266–2301, 2005.
- [11] C.J. Stam and E.C.W. van Straaten, "The organization of physiological brain networks," *Clin. Neurophysiol.*, vol. 123, pp. 1067–1087, 2012.
- [12] B. Tóth, B. File, R. Boha, Z. Kardos, Z. Hidasi, Z.A. Gaál, *et al.*, "EEG network connectivity changes in mild cognitive impairment - Preliminary results," *Int. J. Psychophysiol.*, vol. 92, pp. 1–7, 2014.
- [13] C. Gómez, C. Juan-Cruz, J. Poza, S.J. Ruiz-Gómez, J. Gomez-Pilar, P. Núñez, *et al.*, "Alterations of effective connectivity patterns in mild cognitive impairment: An MEG study," *J. Alzheimer's Dis.* In Press, 2018.
- [14] G. Nolte, A. Ziehe, V.V. Nikulin, A. Schlögl, N. Krämer, T. Brismar, and K.R. Müller, "Robustly estimating the flow direction of information in complex physical systems," *Phys. Rev. Lett.*, vol. 100, pp. 1–5, 2008.
- [15] A. Ewald, F.S. Avarvand, and G. Nolte, "Identifying causal networks of neuronal sources from EEG/MEG data with the phase slope index: a simulation study," *Biomed. Tech. (Berl.)*, vol. 58, pp. 165–178, 2013.
- [16] J. Dauwels, F. Vialatte, F. Musha, and A. Cichocki, "A comparative study of synchrony measures for the early diagnosis of Alzheimer's disease based on EEG," *Neuroimage*, vol. 49, pp. 668–693, 2010.
- [17] Z. Liu, L. Bai, R. Dai, C. Zhong, H. Wang, Y. You, *et al.*, "Exploring the effective connectivity of resting state networks in mild cognitive impairment: an fMRI study combining ICA and mGCA," *Conf. Proc. IEEE Eng. Med. Biol. Soc.*, pp. 5454–5457, 2012.
- [18] F. Vecchio and C. Babiloni, "Direction of information flow in Alzheimer's Disease and MCI patients," *Int. J. Alzheimers. Dis.*, vol. 2011, pp. 1–7, 2011.
- [19] G.B. Karas, P. Scheltens, S.A.R.B. Rombouts, P.J. Visser, R.A. Van Schijndel, N.C. Fox, and F. Barkhof, "Global and local gray matter loss in mild cognitive impairment and Alzheimer's disease," *Neuroimage*, vol. 23, pp. 708–716, 2004.
- [20] C. Gómez, C.J. Stam, R. Hornero, A. Fernández, and F. Maestú, "Disturbed beta band functional connectivity in patients with mild cognitive impairment: an MEG study," *IEEE Trans. Biomed. Eng.*, vol. 56, pp. 1683–1690, 2009.